

9 2610200



©2023 Haleon or its licensed subsidiaries. All trademarks are owned by or licensed to the Haleon group of companies. 0200000201976  
Distributed by: Haleon, Warren, NJ 07059

**TAMPER-EVIDENT BOTTLE**  
DO NOT USE IF INNER SAFETY SEAL UNDER CAP IMPRINTED WITH  
"SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING

**Drug Facts**

**Active ingredients (in each caplet) Purposes**

Acetaminophen 500 mg ..... Pain reliever  
Caffeine 65 mg ..... Pain reliever aid

**Uses**

- temporarily relieves minor aches and pains due to:
  - headache
  - muscular aches

**Warnings**

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.  
**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 6 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much

**Drug Facts (continued)**

caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heart beat.

**Do not use**

- if you are allergic to acetaminophen
- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

**Ask a doctor before use if**

- you have liver disease

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin

**Stop use and ask a doctor if**

- any new symptoms occur
  - painful area is red or swollen
  - pain gets worse or lasts for more than 10 days
  - fever gets worse or lasts for more than 3 days
- These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Drug Facts (continued)**

**Directions**

- do not use more than directed
- adults and children 12 years of age and over: take 2 caplets every 6 hours; not more than 6 caplets in 24 hours
- children under 12 years: ask a doctor

**Other information**

- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- read all product information before using. Keep this box for important information.

**Inactive ingredients** benzoic acid, D&C red no. 27 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C yellow no. 6 aluminum lake, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized corn starch, stearic acid, talc, titanium dioxide

**Questions or comments?** 1-800-468-7746

ASPIRIN-FREE  
Acetaminophen 500 mg, Caffeine 65 mg  
Pain Reliever / Pain Reliever Aid

**EXCEDRIN**  
**TENSION HEADACHE**

HALEON



ACTUAL SIZE



00201976

HALEON

**EXCEDRIN**

**TENSION HEADACHE**

ASPIRIN-FREE

Acetaminophen 500 mg, Caffeine 65 mg  
Pain Reliever / Pain Reliever Aid

ACTUAL SIZE



**24**

**CAPLETS**